Bibliography and further reading

- Appleby L et al, Suicide within 12 months of contact with mental health services: national clinical survey. BMJ, 1999, 318(7193), 1235-1239

- Appleby L et al, National Confidential Inquiry into Suicide and Safety in Mental Health. Annual Report: England, Northern Ireland, Scotland and Wales, Centre for Mental Health and Safety, 2019

- Appleby L et al, National Confidential Inquiry into Suicide and Safety in Mental Health. Annual Report: England, Northern Ireland, Scotland and Wales, Centre for Mental Health and Safety, 2021

- Barrera A et al, Introducing artificial intelligence in acute psychiatric inpatient care: qualitative study of its use to conduct nursing observations, Evidence-based mental health, 2020, 23(1):34-38

- Barnicot K et al, Staff and patient experiences of decision-making about continuous observation in psychiatric hospitals. Social psychiatry and psychiatric epidemiology, 2017, 52(4), 473–483

- Berg SH, Rørtveit K, Walby FA and Aase K, Safe clinical practice for patients hospitalised in mental health wards during a suicidal crisis: qualitative study of patient experiences, BMJ open, 2020, 10(11)

- Berman AL, Risk factors proximate to suicide and suicide risk assessment in the context of denied suicide ideation, Suicide and Life-Threatening Behavior, 2017, 48(3), 340-352.

- Birken M and Bryant W, A Photovoice study of user experiences of an occupational therapy department within an acute inpatient mental health setting, British Journal of Occupational Therapy, 2019, 82(9), 532-543.

- Bowers L et al, The city 128 study of observation and outcomes on acute psychiatric wards, Report to the NHS SDO Programme, December 2006

- Carter G et al, Predicting suicidal behaviours using clinical instruments: systematic review and meta-analysis of positive predictive values for risk scales, The British Journal of Psychiatry, 2017, 210(6), 387-395

- Chambers M et al, The Therapeutic Engagement Questionnaire (TEQ): a service user-focused mental health nursing outcome metric, BMC Psychiatry, 2019; 19: 384

- Chan MK et al, Predicting suicide following self-harm: systematic review of risk factors and risk scales, The British Journal of Psychiatry, 2016, 209(4), 277-283

- Chu S, Special observations in the care of psychiatric inpatients: a review of the literature and developments in practice, ARC Journal of Psychiatry, 2016, 1(1), 21-31

- Department of Health, Best Practice in Managing Risk: Principle and Evidence for Best Practice in Assessment and Management of Risk to Self and Others in Mental Health Services, 2009

- Department of Health, Preventing suicide in England: A cross-government outcomes strategy to save lives, 2011

- Edmondson AC, The fearless organization – Creating Psychological Safety in the Workplace for learning, innovation and growth, Wiley 2019

- Flynn S et al, Suicide by mental health in-patients under observation, Psychological Medicine, 2017 47(13), 2238-2245

- Hall RC Platt DE and Hall RC, Suicide risk assessment: a review of risk factors for suicide in 100 patients who made severe suicide attempts: evaluation of suicide risk in a time of managed care, Psychosomatics, 1999, 40(1), 18-27

- Haney EM et al, Suicide Risk Factors and Risk Assessment Tools: A Systematic Review. Evidence-Based Synthesis Program (ESP) Center, Portland VA Medical Center, Department of Veterans Affairs, March 2012

- Health Education England, Stepping forward to 2020/21 - The mental health workforce plan for england, 2017

- Hunt BR and Lennox BR, Enhancing psychological safety in mental health services, International Journal of Mental Health Systems, 2021, 15(1)

- Hunt IM, Suicide in current psychiatric in-patients: a case-control study, Psychological medicine, 2007, 37(6), 831

- Hunt IM, Windfuhr K, Shaw J, Appleby L and Kapur N, Ligature points and ligature types used by psychiatric inpatients who die by hanging. Crisis, 2012, 33(2), 87-94

- Insua-Summerhays B et al, Staff and patient perspectives on therapeutic engagement during one-to-one observation. Journal of Psychiatric and Mental Health Nursing, 2018, 25(9-10), 546-557

- James K, Stewart D and Bowers L, Self-harm and attempted suicide within inpatient psychiatric services: A review of the literature, International journal of mental health nursing, 2012, 21(4), 301-309

- Jovanović N, Campbell J and Priebe S, How to design psychiatric facilities to foster positive social interaction – a systematic review. Eur Psychiatry, 2019, 60, 49-62

- Kapur N, Gorman LS, Quinlivan L and Webb RT, Mental health services: quality, safety and suicide, BMJ Quality and Safety, 2021, 31(6)

- Levi-Belz Y et al, Psychosocial factors correlated with undisclosed suicide attempts to significant others: findings from the adolescence SEYLE study, Suicide and Life-Threatening Behavior, 2019, 49(3), 759-773

- Lyubovnikova J, West MA, 'Why teamwork matters: enabling health care team effectiveness for the delivery of high quality patient care' in Salas S, Tannembaum I, Cohen D, Latham G (eds), Developing and enhancing teamwork in organisations, 2013, p331–72

- Madsen T, Erlangsen A and Nordentoft M, Risk estimates and risk factors related to psychiatric inpatient suicide – An overview, International journal of environmental research and public health, 2017 14(3), 253

- Nanda U, Eisen S, Zadeh RS and Owen D, Effect of visual art on patient anxiety and agitation in a mental health facility and implications for the business case, Journal of Psychiatric and Mental Health Nursing, 2011, 18(5), 386-393

- National Confidential Inquiry into Suicide and Homicide by People with Mental Illness, Healthy Services & Safer Patients: links between patient suicide and mental health care providers, February 2015

- National Confidential Inquiry into Suicide and Safety in Mental Health, Safer Services: A toolkit for specialist mental health services and primary care, 2021

- NHS England, Culture and Values for Workplace Compassion, 2018

- NHS England and Health Education England, Self harm and suicide prevention competence framework – Adults and older adults, 2018

- National Quality Board, Supporting NHS Providers to Deliver the Right Staff, with the Right Skills, in the Right Place at the Right Time: Safe, Sustainable and Productive Staffing, NHS England, 2016

- Oostermeijer S et al, Design features that reduce the use of seclusion and restraint in mental health facilities: a rapid systematic review, BMJ Open, 2021 11(7)

- Quinlivan L et al, Which are the most useful scales for predicting repeat self-harm? A systematic review evaluating risk scales using measures of diagnostic accuracy. BMJ Open, 2016, 6(2)

- Rahman M et al, Nile Ward PICU violence reduction quality improvement project, BJPsych Open, 2021, 7(S1), S213-S214

- Sakinofsky I, Preventing suicide among inpatients, Can J Psychiatry, 2014, 59(3), 131-140

- Thomas JB and Haslam CO. [How people who self-harm negotiate the inpatient environment: the mental healthcare workers perspective,](#) J Psychiatr Ment Health Nurs. 2017, 24(7):480-490

- Turner, K., Stapelberg, N. J., Sveticic, J., & Dekker, S. W. (2020). [Inconvenient truths in suicide prevention: Why a Restorative Just Culture should be implemented alongside a Zero Suicide Framework](#). Australian & New Zealand Journal of Psychiatry, 54(6), 571-581

- Ulrich RS, [Visual landscapes and psychological wellbeing](#), Landscape research, 1979, 4(1), 17-23

- Van der Schaaf PS et al, [Impact of the physical environment of psychiatric wards on the use of seclusion](#), *BJPsych*, 2013, *202*(2), 142-149

- Wand T, [Considering the culture of blame in mental health care and service delivery,](#) International Journal of Mental Health Nursing, 2017, 26: 3–4.

- Wyder M et al, [How health care systems let our patients down: a systematic review into suicide deaths](#), International Journal for Quality in Health Care, June 2020, 32(5), 285-291

- Walsh G, Sara G, Ryan CJ and Large M, [Meta-analysis of suicide rates among psychiatric inpatients](#), Acta Psychiatrica Scandinavica, 2015 131(3), 174-184